

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 September 2001 (07.09.2001)

(10) International Publication Number
PCT WO 01/64645 A2

(51) International Patent Classification⁷: C07D 215/00

FIN-20100 Turku (FI). SALO, Harri [FI/FI]; Puolikontie 6 C, FIN-20300 Turku (FI).

(21) International Application Number: PCT/FI01/00203

(22) International Filing Date: 28 February 2001 (28.02.2001)

(74) Agent: ORION CORPORATION; ORION PHARMA, Industrial Property Rights, P.O.Box 65, FIN-02101 Espoo (FI).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
20000480 1 March 2000 (01.03.2000) FI

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicants (*for all designated States except US*): ORION CORPORATION [FI/FI]; Orionintie 1, FIN-02200 Espoo (FI). OY JUVANTIA PHARMA LTD [FI/FI]; Tykistökatu 6 A, BioCity, FIN-20520 Turku (FI).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): WURSTER, Siegfried [DE/FI]; Matarakaari 28 A 1, FIN-21500 Piiikkiö (FI). ENGSTRÖM, Mia [FI/FI]; Sirkkalankatu 12 A 6, FIN-20500 Turku (FI). SAVOLA, Juha-Matti [FI/FI]; Siutlankatu 8, FIN-20380 Turku (FI). HÖGLUND, Iisa [FI/FI]; Kaniikintie 17, FIN-20300 Turku (FI). SALLINEN, Jukka [FI/FI]; Puutarhakatu 24 C 32, FIN-20100 Turku (FI). HAAPALINNA, Antti [FI/FI]; Markulantie 8 A, FIN-20360 Turku (FI). TAUBER, Andrei [RU/FI]; Väärämäentie 7 d 154, FIN-00700, Helsinki (FI). HOF-FRÉN, Anna-Marja [FI/FI]; Bilmarkinkatu 4 as 12,

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DERIVATIVES OF QUINOLINE AS ALPHA-2 ANTAGONISTS



(57) Abstract: The invention provides a compound of formula (I), wherein A, Ra, Rb, R₁ to R₅, m and t are as defined as disclosed, or a pharmaceutically acceptable salt or ester thereof, useful as alpha-2 antagonist. The compounds I can be used for the treatment of diseases or conditions wherein alpha-2 antagonists are indicated to be effective.

THIS PAGE BLANK (USPTO)

DERIVATIVES OF QUINOLINE AS ALPHA-2 ANTAGONISTS

FIELD OF THE INVENTION

The present invention relates to therapeutically active derivatives of quinoline, including the pharmaceutically acceptable salts and esters thereof,
5 and their use as alpha-2 antagonists.

BACKGROUND

Some compounds exhibiting alpha adrenergic activity are well known in the art. Those compounds may be used for the treatment of a wide variety of diseases and conditions of the peripheric system and the central nervous
10 system (CNS).

The alpha adrenergic receptors are divided into alpha-1 and alpha-2 adrenoceptors, each of which are further divided into subtypes. Accordingly, alpha-2 adrenoceptors in humans have been subdivided into three pharmacological subtypes known as alpha-2A, alpha-2B and alpha-2C
15 adrenoceptors. A fourth subtype, alpha-2D, is known in rat, bovine and porcine and it corresponds to alpha-2A in man. These subtypes have a distinct distribution in human and animal tissues. For instance, alpha-2C adrenoceptors are concentrated in the CNS and they appear to play a role in the modulation of various CNS-mediated behavioural and physiological responses.

20 Compounds that are non-specific to any of the above-mentioned alpha-2 subtypes, and compounds that are specific to a certain alpha-2 subtypes are already known. For example, atipamezole is a non-specific alpha-2 antagonists. Atipamezole has been described in, for example, EP-A-183 492 (cf. p.13, compound XV) and A. Haapalinna et al., Naunyn-Schimiedeberg's
25 Arch. Pharmacol. Vol. 356, 1997, p.570-582. U.S. Patent No. 5,902,807 describes compounds that are selective antagonists for the alpha-2C subtype and may be used in the treatment of mental illnesses, e.g. mental disturbances induced by stress. Such compounds include, for example, MK-912 and BAM-1303. Furthermore, WO-A-99 28300 discloses substituted imidazole derivatives
30 having agonist-like activity for alpha-2B or 2B/2C adrenoceptors. The disclosures of all documents cited above in this paragraph are incorporated by reference herein.

As to the derivatives of quinoline, Medicinskaja parazitologija I parazitarnye bolezni, vol.5, 1991, 55-7 (Mikhailitsyn F.S. et al.) and J. Med. Chem., vol.20(8), 1977, 987-996 (Cain F.C. et al.) describe, for example, acridine derivatives as anticancer and/or antiparasitic agents. In addition, a publication by Adams et al in 1985 (Mol. Pharm. 27, 480-491) reports on the binding of diquinolines, diacridines and a number of monoacridines on rat brain alpha-1-, alpha-2- and beta-adrenoceptors.

SUMMARY OF THE INVENTION

An object of the present invention is to provide further antagonists of alpha-2 adrenoceptors that can be used for the treatment of diseases or conditions of the peripheral or central nervous system where alpha-2 antagonists are indicated to be useful. Accordingly, an object of the present invention is to provide further compounds to be used as alpha-2 antagonist agents in the treatment of mammals, including humans and animals.

Another object of the present invention is to provide further compounds useful as selective alpha-2C antagonist agents for the treatment of various disorders or conditions of the central nervous system where alpha-2C antagonists are indicated to be useful.


BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the effects of atipamezole and compound 1 on dexmedetomidine-induced hypolocomotion.

Figure 2 shows the effects of atipamezole and compound 1 in a forced swimming test in Balb/c mice.

DETAILED DESCRIPTION OF THE INVENTION

One embodiment of the present invention covers compounds of formula I:



wherein,

5 R_1 is H or (C₁-C₆)alkyl;

each R_2 is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or hydroxy(C₁-C₆)alkyl;

10

A is a benzene ring or (C₅-C₇)cycloalkyl;


when A is a benzene ring each R_3 is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbonyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

15

when A is (C₅-C₇)cycloalkyl each R_3 is independently OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;

20

R_4 and R_5 form, together with the N-atom to which they are attached,




25

wherein X is O or =NR₆; R_6 is H, OH, NH₂, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, CN-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-

4

CO-, NH₂-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, phenyl, naphthyl or benzyl, wherein the said phenyl, naphthyl or benzyl is optionally substituted with one to three substituent(s) each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, 5 mono- or di(C₁-C₆)alkylamino or halo-(C₁-C₆)alkyl;

or R₄ and R₅ form, together with the N-atom to which they are attached,



10 wherein n = 1 or 2; R₆ is as defined above; and r = 0 to 3;

or R₄ and R₅ form, together with the N-atom to which they are attached, 1-imidazolyl, 1-imidazolynyl or 1-triazolyl, each of which can optionally be substituted with one to three substituent(s) R₇ each independently being (C₁-C₆)alkyl or NH₂; 15

or one of R₄ and R₅ is -SO₂R₈ and the other of R₄ and R₅ is H or (C₁-C₆)alkyl; R₈ is (C₁-C₆)alkyl, phenyl, naphthyl or benzyl, wherein the said phenyl, naphthyl or benzyl is optionally substituted with one to three substituent(s) R₉ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino; 20

R_a and R_b are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, 25 mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN;

or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed benzene ring optionally substituted with one to three substituent(s) R'₃ each independently being OH, halogen, (C₁-C₆)alkyl, 30 (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

5

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally substituted with one to four substituent(s) R₁₀ each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or
5 hydroxy(C₁-C₆)alkyl;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed bicyclo[2.2.1]-heptane ring optionally substituted with one to four substituent(s) each independently being OH, halogen, (C₁-
10 C₆)alkyl or (C₁-C₆)alkoxy;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five or six membered heterocyclic ring with one ring heteroatom =NR₁₁, which heterocyclic ring is optionally substituted with one to
15 three substituent(s) R₁₀ as defined above; R₁₁ is H or (C₁-C₆)alkyl, or R₁₁ is phenyl optionally substituted with one to three substituents R₁₂ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino;

20 m is 0 to 3; and
t is 0 to 3,

or pharmaceutically acceptable salts and esters thereof.

The compounds of formula I can be used for the manufacture of a
25 medicament for the treatment of diseases or conditions where alpha-2 antagonists are indicated to be effective.

The following subgroups (1) to (18) of compounds of formula I taken alone or in any combination with each other are possible:

- 30 (1) A is a benzene ring;
(2) A is a (C₅-C₇)cycloalkyl;
(3) Ra and Rb are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN; e.g. H,

6

halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy or halo(C₁-C₆)alkyl; such as H, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy; e.g. H or (C₁-C₃)alkyl, wherein the said (C₁-C₃)alkyl includes both straight and branched chain radicals of up to 3 carbon atoms;

5

(4) Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed benzene ring unsubstituted or substituted with one to three, e.g. one or two, such as one, substituent(s) R'₃ each independently being OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, NH₂-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, (C₁-C₆)alkyl-S or hydroxy(C₁-C₆)alkyl; e.g. OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl or (C₁-C₆)alkoxy; such as halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy; e.g. (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

10

15

(5) Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally substituted with one to four, e.g. one or two, such as one, substituent(s) R₁₀ each independently being OH, halogen, (C₁-C₆)alkyl (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl; such as (C₁-C₆)alkyl or (C₁-C₆)alkoxy; e.g. (C₁-C₆)alkyl;

20

(6) Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed bicyclo[2.2.1]-heptane ring optionally substituted with one to four, e.g. one or two, such as one, substituent(s) each independently being OH, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy; e.g. (C₁-C₆)alkyl

25

(7) Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five or six membered heterocyclic ring with one ring heteroatom =NR₁₁, which heterocyclic ring is optionally substituted with one to three, e.g. one or two, such as one substituent(s) R₁₀ as defined above; and R₁₁ is H or (C₁-C₆)alkyl, or R₁₁ is phenyl optionally substituted with one to three, e.g. one or two, such as one, substituents R₁₂ each independently being OH,


30

halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino;

- 5 (8) R₄ and R₅ form, together with the N-atom, 1-piperazinyI which is 4-substituted with R₆, wherein R₆ is as defined above; e.g. (C₁-C₆)alkyl, (C₂-C₆)alkenyl, CN-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-, NH₂-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl or optionally substituted benzyl; such as (C₁-C₆)alkyl, (C₂-C₆)alkenyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl; e.g. (C₁-C₆)alkyl;
- 10 (9) R₄ and R₅ form, together with the N-atom, a morpholino ring;
- (10) R₄ and R₅ form, together with the N-atom, a 1-piperidinyI or 1-pyrrolidinyI optionally substituted with R₆, wherein R₆ is as defined above, possibly (C₁-C₆)alkyl, (C₂-C₆)alkenyl, CN-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-, NH₂-CO-, mono- or
15 di(C₁-C₆)alkylcarbamoyl, hydroxy(C₁-C₆)alkyl or (C₃-C₆)cycloalkyl; such as (C₁-C₆)alkyl, (C₂-C₆)alkenyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl; e.g. (C₁-C₆)alkyl;
- (11) R₄ and R₅ form, together with the N-atom, 1-imidazolyl, 1-imidazolinyI or 1-triazolyl, each of which can optionally be substituted
20 with one to three, e.g. one or two, such as one, substituent(s) R₇ each independently being (C₁-C₆)alkyl or NH₂; e.g. R₄ and R₅ form, together with the N-atom, 2-amino-imidazol-1-yl or 2-amino-imidazolin-1-yl;
- (12) one of R₄ and R₅ is -SO₂R₈ and the other of R₄ and R₅ is H or
25 (C₁-C₆)alkyl; R₈ is independently (C₁-C₆)alkyl, phenyl, naphthyl and benzyl, wherein the said phenyl, naphthyl or benzyl is optionally substituted with one to three, e.g. one or two, such as one, substituent(s) R₉ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino;
- 30 (13) m is 0 to 3; e.g. 0, 1 or 2; e.g. 0 or 1; such as 0;
- (14) t is 0 to 3; e.g. 0, 1 or 2; e.g. 0 or 1; such as 0; and/or
- (15) R₁ is H;

- (16) R_1 is (C_1-C_6) alkyl;
- (17) R_2 is independently OH, halogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_1-C_6) alkoxy, halo (C_1-C_6) alkyl, NO_2 , NH_2 , mono- or di (C_1-C_6) alkylamino, (C_1-C_6) alkyl-S- or hydroxy (C_1-C_6) alkyl; such as OH, halogen, (C_1-C_6) alkyl, (C_1-C_6) alkoxy or hydroxy (C_1-C_6) alkyl; e.g. (C_1-C_6) alkyl or (C_1-C_6) alkoxy; and/or
- (18) R_3 is independently OH, halogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_1-C_6) alkoxy, halo (C_1-C_6) alkyl, NO_2 , NH_2 , mono- or di (C_1-C_6) alkylamino, (C_1-C_6) alkyl-CO-, mono- or di (C_1-C_6) alkylcarbamoyl, (C_1-C_6) alkyl-S-, hydroxy (C_1-C_6) alkyl or NH_2 -CO-; such as OH, halogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_1-C_6) alkoxy, halo (C_1-C_6) alkyl; e.g. halogen, (C_1-C_6) alkyl or (C_1-C_6) alkoxy.


A possible subgroup of the compound of formula I is a compound of formula IA:



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} ; m and t are as defined above; i is 1 to 3; and j is 0 to 4.


9

Another possible subgroup of the compound of formula I is a compound of formula IB:



- wherein A, R₁, R₂, R₃, R₄, R₅, Ra, Rb, m and t are as defined above;
- 5 and Ra and Rb are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN; or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed
- 10 five to seven membered carbocyclic ring optionally substituted with one to three substituent(s) R₁₀; each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl.


Another possible subgroup of the compound of formula I is a compound of formula IC:



10


wherein $R_1, R_2, R_3, R_4, R_5, R_{10}, R_{11}, m$ and t are as defined above; i is 1 or 2; and j is 0 to 3.

Another possible subgroup of the compound of formula I is a compound
5 of formula ID:




wherein $R_1, R_2, R_3, R'_3, R_4, R_5, m$ and t are as defined above and p is 0 to 3; for example wherein m is 1 and R_3 is (C_1-C_6) alkoxy.

10 In a possible subgroup of the compound of formula I, IA, IB, IC or ID, R_4 and R_5 form, together with the N-atom to which they are attached,



wherein X is $=NR_6$; R_6 is as defined above under formula I; such as (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy (C_1-C_6) alkyl or (C_3-C_6) cycloalkyl; e.g. (C_1-C_6) alkyl.
15

Another embodiment of the invention provides new compounds of formula II:



wherein,

5

R_1 is H or (C₁-C₆)alkyl;

each R_2 is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or hydroxy(C₁-C₆)alkyl;

10

A is a benzene ring or (C₅-C₇)cycloalkyl;


when A is a benzene ring each R_3 is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

15

when A is (C₅-C₇)cycloalkyl each R_3 is independently OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;

20

R_4 and R_5 form, together with the N-atom to which they are attached,




wherein X is O or =NR₆; R_6 is H, OH, NH₂, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, CN-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-, NH₂-CO-,

25

mono- or di(C₁-C₆)alkylcarbamoyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl or benzyl, wherein the said benzyl is optionally substituted with one to three substituent(s) each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or halo-(C₁-C₆)alkyl;

5

or R₄ and R₅ form, together with the N-atom to which they are attached,



wherein n = 1 or 2; R₆ is as defined above; and r = 0 to 3;

10 or R₄ and R₅ form, together with the N-atom to which they are attached, 1-imidazolyl, 1-imidazoliny or 1-triazolyl, each of which can optionally be substituted with one to three substituent(s) R₇ each independently being (C₁-C₆)alkyl or NH₂;

15 R_a and R_b are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN;

20 or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed benzene ring optionally substituted with one to three substituent(s) R'₃ each independently being OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

25

or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally substituted with one to four substituent(s) R₁₀ each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;

30

or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed bicyclo[2.2.1]-heptane ring optionally substituted with

one to four substituent(s) each independently being OH, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

or Ra and Rb form, together with the carbon ring atoms to which they are
5 attached, a condensed five or six membered heterocyclic ring with one ring heteroatom =NR₁₁, which heterocyclic ring is optionally substituted with one to three substituent(s) R₁₀ as defined above; R₁₁ is H or (C₁-C₆)alkyl, or R₁₁ is phenyl optionally substituted with one to three substituents R₁₂ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or
10 mono- or di(C₁-C₆)alkylamino;

m is 0 to 3; and

t is 0 to 3,

15 or of a pharmaceutically acceptable salt or ester thereof, with the provisos, that

a) when A is a benzene ring, m is 0 or 1, t is 0, R₁ is H, R₃ is Cl or NO₂, and Ra and Rb form, together with the carbon ring atoms to which they are
20 attached, a condensed benzene ring, and X is NR₆, then R₆ is not H, -CH₃, -CH₂CH₃, -COCH₃, or -CO-NH₂;


b) when A is a benzene ring, then Ra and Rb are not at the same time H;

c) when A is a benzene ring, m is 1, t is 0, R₁ is H, and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed
25 benzene ring, which is optionally substituted with Br, and X is O, then R₃ is not NO₂ or -OCH₃;

d) when A is a benzene ring, m is 0, t is 0, R₁ is H, and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed nonsubstituted benzene ring, then X is not O;

30 e) the compound is not 4-[4-[(7-Chloro-2-methyl-4-quinoliny)amino]phenyl]-1-diethylcarbamoylpiperazine, 4-[4-[(6-Chloro-2-methoxy-9-acridiny)amino]phenyl]-1-diethylcarbamoylpiperazine, 6-amino-4-[[3-chloro-4-(1H-imidazol-1-yl)phenyl]amino]-7-methoxy-3-quinolinecarbonitrile or 4-[[3-chloro-4-(1H-imidazol-1-yl)phenyl]amino]-7-methoxy-6-nitro-3-
35 quinolinecarbonitrile.

A possible subgroup of the compound of formula II is a compound of formula IIA,



5 wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} , m and t are as defined in formula II; i is 1 to 3; and j is 0 to 4; for example

 wherein i is 2, j is 0 or 1 and R_{10} is (C_1-C_3) alkyl, wherein the (C_1-C_3) alkyl includes both straight and branched chain radicals of up to 3 carbon atoms; or

10


 wherein m is 0 or 1 and R_3 is (C_1-C_3) alkyl or halogen, wherein the (C_1-C_3) alkyl includes both straight and branched chain radicals of up to 3 carbon atoms; or

15

 wherein the compound is [4-(4-Methylpiperazin-1-yl)phenyl]-(1,2,3,4-tetrahydroacridin-9-yl)amine, 2-{4-[4-(1,2,3,4-Tetrahydroacridin-9-yl)aminophenyl]piperazin-1-yl}ethanol, [4-(4-Methylpiperazin-1-yl)phenyl]-(2-methyl-1,2,3,4-tetrahydro-acridin-9-yl)amine, (8-Fluoro-1,2,3,4-tetrahydroacridin-9-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, [4-(4-Methylpiperazin-1-yl)phenyl]-(2,7-dimethyl-1,2,3,4-tetrahydroacridin-9-yl)amine, [4-(4-Methylpiperazin-1-yl)phenyl]-(7,8,9,10-tetrahydro-6H-cyclohepta[b]quinolin-11-yl)amine or [4-(4-Methylpiperazin-1-yl)phenyl]-(1,1,3,3-tetramethyl-1,2,3,4-tetrahydroacridin-9-yl)amine.

25

 Another possible subgroup of the compound of formula II is a compound of formula IIB,



wherein A, R₁, R₂, R₃, R₄, R₅, m and t are as defined in formula II; and

Ra and Rb are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂,
 5 mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN; or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally substituted with one to three
 10 substituent(s) R₁₀; each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl; for example

wherein A is a benzene ring; or

wherein m is 0 or 1; or

15

wherein R₃ is (C₁-C₆)alkyl or (C₁-C₆)alkoxy; or


wherein Ra and Rb are independently H or (C₁-C₃)alkyl, wherein the (C₁-C₃)alkyl includes both straight and branched chain radicals of up to 3 carbon
 20 atoms; or

wherein A is a six membered carbocyclic ring and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring, wherein the carbocyclic ring can
 25 optionally be substituted with one to three substituent(s) each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or hydroxy(C₁-C₆)alkyl; or

16

wherein the compound is (3-Ethyl-2,8-dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (2-Methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (3-Ethyl-2,6-dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (3-Ethyl-6-methoxy-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (3-Ethyl-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, 4-[4-(4-Methylpiperazin-1-yl)phenylamino]quinoline-3-carbonitrile, 3-Isopropyl-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (2,3-Dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, [4-(4-Methylpiperazin-1-yl)phenyl]-(1,2,3,4,5,6,7,8-octahydroacridin-9-yl)amine or (3-Ethyl-2-methylquinolin-4-yl)-methyl-[4-(4-methylpiperazin-1-yl)phenyl]amine.


Another possible subgroup of the compound of formula II is a compound of formula IIC,



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} , R_{11} , m and t are as defined in formula II; i is 1 or 2; and j is 0 to 3.

20

Another possible subgroup of the compound of formula II is a compound of formula IID,



wherein R_1 , R_2 , R_3 , R'_3 , R_4 , R_5 , m and t are as defined in formula II; and p is 0 to 3; for example


wherein m is 1 and R_3 is (C_1-C_6) alkoxy; or

5

wherein the compound is 2-{4-[4-(Acridin-9-yl)aminophenyl]piperazin-1-yl}-ethanol, (4-Methoxyacridin-9-yl)-[4-(4-methylpiperazin-1-yl)-phenyl]amine, Acridin-9-yl-[4-(piperidin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-benzylpiperazin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-methylpiperidin-1-yl)phenyl]amine, Acridin-9-yl-[4-(3-hydroxymethylpiperidin-1-yl)phenyl]amine, Acridin-9-yl-[4-pyrrolidin-1-yl]phenyl]amine, Acridin-9-yl-[4-(4-cyclopropylpiperazin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-isopropylpiperazine-1-yl)phenyl]amine, (Acridin-9-yl)-methyl-[4-(4-methylpiperazin-1-yl)phenyl]amine or Acridin-9-yl-[2,5-diethoxy-4-(morpholin-4-yl)phenyl]amine.

15

In a possible subgroup of the compound of Formula II, IIA, IIB, IIC, or IID, R_4 and R_5 form, together with the N-atom to which they are attached,



20

wherein R_6 is (C_1-C_6) alkyl, (C_1-C_6) alkenyl, (C_3-C_6) cycloalkyl or hydroxy (C_1-C_6) alkyl, for example, R_6 is (C_1-C_6) alkyl.

The compounds of formulae I and II and the subgroups IA, IB, IC, ID, IIA, IIB, IIC, and IID thereof, as well as the pharmaceutically acceptable esters

and salts thereof, are referred to below as the compounds of the invention, unless otherwise indicated.

5 The compounds of the invention may have chiral carbon atom(s) in their structure. The invention includes within its scope all the possible stereoisomers of the compounds, including geometric isomers, e.g. Z and E isomers (cis and trans isomers), and optical isomers, e.g. diastereomers and enantiomers. Furthermore, the invention includes in its scope both the individual isomers and any mixtures thereof, e.g. racemic mixtures. The individual isomers may be
10 obtained using the corresponding isomeric forms of the starting material or they may be separated after the preparation of the end compound according to conventional separation methods. For the separation of, for example, optical isomers, e.g. enantiomers, from the mixture thereof the conventional resolution methods, e.g. fractional crystallisation, may be used.

15 Physiologically acceptable salts, e.g. acid addition salts with both organic and inorganic acids are well known in the field of pharmaceuticals. Non-limiting examples of these salts include chlorides, bromides, sulfates, nitrates, phosphates, sulfonates, formates, tartrates, maleates, citrates, benzoates, salicylates and ascorbates. Pharmaceutically acceptable esters, when
20 applicable, may be prepared by known methods using pharmaceutically acceptable acids that are conventional in the field of pharmaceuticals and that retain the pharmacological properties of the free form. Non-limiting examples of those esters include esters of aliphatic or aromatic alcohols, e.g. lower alkyl esters, e.g. methyl, ethyl and propyl esters.


25 Terms employed herein have the following meanings: A halogen or halo refers to fluorine, chlorine, bromine or iodine, e.g. fluorine or chlorine. The term (C₁-C₆)alkyl as employed herein as such or as part of another group includes both straight, and branched chain radicals of up to 6 carbon atoms, for example of 1 to 4 carbon atoms, e.g. methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl or s-butyl. The term (C₁-C₆)alkoxy as such or as part of another group refers to -O-
30 (C₁-C₆)alkyl, wherein (C₁-C₆)alkyl is as defined above. The term (C₂-C₆)alkenyl includes both straight and branched chain radicals of up to 6 carbon atoms, for example of 2 to 4 carbon atoms, containing (a) double bond(s). The term (C₂-C₆)alkynyl includes both straight and branched chain radicals of up to
35 6 carbon atoms, for example of 2 to 4 carbon atoms, containing (a) triple

bond(s). The term halo-(C₁-C₆)alkyl refers to (C₁-C₆)alkyl radical, as defined above, that is substituted by one or more halo radicals as defined above, for example trifluoromethyl, difluoromethyl etc. The term mono- or di(C₁-C₆)alkylcarbamoyl refers to a carbamoyl radical which is N-substituted with one or two (C₁-C₆)alkyl radical(s), as defined above.

The compounds of the invention can be prepared analogously or according to a variety of synthetic routes known in the literature using suitable starting materials, for example, according to or analogous to methods described by B.F. Cain et al. in J.Med.Chem., vol.20(8), 1977, pp. 987-996, and by W.A. Denny et al. in J.Med.Chem., vol.25, 1982, p.276-315, the contents of which are hereby incorporated by reference.

In general, the compounds of the invention can be prepared e.g. analogously or according to the following reaction scheme 1:

Scheme 1



15

wherein Ra, Rb, R₁, R₂, R₃, R₄, R₅, m and t are as defined above.

The above reaction is a conventional acid-catalyzed coupling of the chloro-compound of formula III with a substituted aromatic amine of formula IV. The reaction is carried out at room or elevated temperature in a suitable solvent, e.g. alcohol, such as methanol, in acidic conditions, to obtain the compound of formula I' which is then isolated from the reaction mixture in a usual manner.


The starting compounds III and IV are commercially available or can be prepared according to or analogously to the methods described in the literature

(see e.g. J.Med.Chem., vol.20(8), 1977, pp. 987-996, and J.Med.Chem., vol.25, 1982, p.276-315, as mentioned above).

Accordingly, a substituted aromatic NR₁-amine IV can e.g. be prepared starting from the corresponding nitro compound which is reduced with a reducing agent, e.g. in the presence of SnCl₂·H₂O, in a suitable solvent, e.g. DMF, and optionally alkylated with R₁ in a manner known in the art, when R₁ = (C₁₋₆)alkyl is desired.

The starting material III can be prepared e.g. according to following scheme 2:

10 Scheme 2



wherein Ra, Rb, R₃, and m are as defined above.

In scheme 2 a compound V is reacted e.g. with thionyl chloride in the presence of a small amount of DMF, to obtain the 9-chlorinated reactant III.
15 The reaction is carried out at room or elevated temperature.

It is obvious for a skilled person that, in the above reactions, any starting material or intermediate can be protected, if necessary, in a manner well known in the chemical field. Any protected functionality can subsequently be deprotected in a manner known in the art.

20 The above disclosed synthetic routes are meant to illustrate the preparation of the compounds of the invention and the preparation is by no means limited thereto, i.e. other synthetic methods that are within the general knowledge of a skilled person are also possible.

25 The compounds of the invention may be converted, if desired, into their pharmaceutically acceptable salt or ester form using methods well known in the art.

The following examples are meant only for illustrating purposes and does not limit the scope of the invention defined in claims.

EXAMPLE 1

5 **Acridin-9-yl-[4-(4-methylpiperazin-1-yl)-phenyl]amine**

Step 1

1.04 g (5.0 mmol) of N-(4-nitrophenyl)piperazine was dissolved in 5 ml of dimethylformamide. Sodium hydride (0.24 g, 6.0 mmol) was added to the reaction mixture in three portions under nitrogen atmosphere and cooling over
10 a period of 10 min. After 30 min of stirring, 0.31 ml (6.0 mmol) of methyl iodide was added dropwise to the reaction mixture at 0 °C. Stirring was continued for 1 h at room temperature, the reaction mixture was then evaporated to dryness, and purified by silica gel chromatography (methylene chloride : methanol : triethylamine 94 : 5 : 1) to yield 1.0 g (90 %) of 1-methyl-4-(4-
15 nitrophenyl)piperazine.

Step 2

0.999 g (4.5 mmol) of 1-methyl-4-(4-nitrophenyl)piperazine, 10.15 g (45 mmol) of tin(II)chloride dihydrate and 20 ml of dimethylformamide were mixed and stirred overnight at 80 °C. Most of dimethylformamide was evaporated
20 under vacuum. The remaining slurry was poured into ice water, neutralized with a sodium bicarbonate (sat.) solution, and filtered. The filtrate was extracted several times with ethylacetate and chloroform to provide 0.636 g (74 %) of 4-(4-methylpiperazin-1-yl)aniline.

Step 3

0.488 g (2.5 mmol) of 9-(10H)acridone, 2.5 ml of thionyl chloride and a catalytic amount (a few drops) of dimethylformamide were mixed at 80 °C. After 30 min of stirring, the reaction mixture was evaporated, the residue was dissolved in chloroform and poured into cold aqueous ammonia. The ammonia solution was extracted several times with chloroform. The combined organic
25 phases were washed with 2M ammonia solution, dried over sodium sulfate and
30 evaporated to obtain 0.517 g (97 %) of 9-chloroacridine.

Step 4

0.191 g (1.0 mmol) of 4-(4-methylpiperazin-1-yl)aniline, 2.5 ml of methanol and a few drops of concentrated hydrochloric acid were mixed and heated under reflux. The 9-chloroacridine (1-1,5 equivalents) and 2.5 ml
5 methanol were mixed separately and added to the reaction mixture in small portions. After 30 min of stirring, two drops of concentrated hydrochloric acid were added, and heating was continued for 2 h. The reaction mixture was then evaporated to dryness and purified by chromatography (silica gel column; gradient from 100 % methylene chloride to 90 % methylene chloride and 10 %
10 methanol; when 4-(4-methylpiperazin-1-yl)aniline eluted from the column, the eluent was changed to methylene chloride : methanol : triethylamine 94 : 5 : 1). A final amount of 0.126 g (34 %, overall yield 12 %) of the title compound in pure form was obtained.

¹H NMR (DMSO-d₆, 500 MHz): 8.05 (2H, m), 7.73 (2H, m), 7.66 (2H, m),
15 7.13 (2H, m), 6.97 (4H, m), 3.35 (4H, m), 2.98 (4H, m), 2.58 (3H, s); MS (EI⁺): 368 (M⁺)

EXAMPLE 2

Acridin-9-yl-[2,5-diethoxy-4-(morpholin-4-yl)phenyl]amine

20 Following the procedure outlined in Step 4 of Example 1, but substituting 2,5-diethoxy-4-morpholinoaniline dihydrochloride for 4-(4-methylpiperazin-1-yl)aniline, afforded the title compound with an overall yield of 23 %.

¹H NMR (CDCl₃, 500 MHz): 8.19 (2H, m), 8.13 (2H, m), 7.56 (1H, s),
7.42 (2H, m), 7.04 (2H, m), 6.42 (1H, s), 3.91 (4H, m), 3.15 (4H, m), 3.10 (4H,
25 q, J = 7.27 Hz), 1.41 (6H, t, J = 7.27 Hz); MS (ESI⁺ TOF): 444 (M⁺)

EXAMPLE 3

Acridin-9-yl-[4-(morpholin-4-yl)phenyl]amine

Following the procedure outlined in Step 4 of Example 1, but substituting
30 4-(morpholin-1-yl)aniline for 4-(4-methylpiperazin-1-yl)aniline, afforded the title compound with an overall yield of 64 %.

23

¹H NMR (CDCl₃, 500 MHz): 7.99 (4H, m), 7.55 (2H, m), 7.17 (2H, m), 7.04 (2H, m), 6.88 (2H, m), 3.88 (4H, m), 3.15 (4H, m); MS (ESI⁺ TOF): 356 (M⁺)

5 **EXAMPLE 4**

(3-Ethyl-2,8-dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure outlined in Step 4 of Example 1, but substituting 4-chloro-2,8-dimethyl-3-ethylquinoline for 9-chloroacridine, afforded the title compound with 27 % overall yield.

¹H NMR (CDCl₃, 500 MHz): 7.60 (1H, m), 7.41 (1H, m), 7.16 (1H, m), 6.79 (2H, m), 6.62 (2H, m), 5.62 (1H, s), 3.12 (4H, m), 2.79 (3H, s), 2.79 (2H, q, J = 7.63 Hz), 2.78 (3H, s), 2.59 (4H, m), 2.36 (3H, s), 1.15 (3H, t, J = 7.63 Hz); MS (ESI⁺ TOF): 375 (M⁺)

15

EXAMPLE 5

(2-Methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure outlined in Step 4 of Example 1, but substituting 4-chloro-2-methylquinoline for 9-chloroacridine, afforded the title compound with 24 % overall yield.

¹H NMR (CDCl₃, 500 MHz): 8.02 (1H, m), 7.91 (1H, m), 7.64 (1H, m), 7.44 (1H, m), 7.23 (2H, m), 7.00 (2H, m), 6.59 (1H, m), 3.26 (4H, m), 2.61 (4H, m), 2.55 (3H, s), 2.38 (3H, s); MS (ESI⁺ TOF): 333 (M⁺)

25 **EXAMPLE 6**

[4-(4-Methylpiperazin-1-yl)phenyl]-(quinolin-4-yl)amine

Following the procedure outlined in Step 4 of Example 1, but substituting 4-chloroquinoline for 9-chloroacridine, afforded the title compound with 11 % overall yield.

24

¹H NMR (CDCl₃, 500 MHz): 8.28 (1H, m), 8.25 (1H, m), 8.06 (1H, m), 7.66 (1H, m), 7.50 (1H, m), 7.25 (2H, m), 6.97 (2H, m), 3.25 (4H, m), 2.64 (4H, m), 2.40 (3H, s); MS (ESI⁺ TOF): 319 (M⁺)

5 **EXAMPLE 7**

(3-Ethyl-2,6-dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure outlined in Step 4 of Example 1, but substituting 4-chloro-1,6-dimethyl-3-ethylquinoline for 9-chloroacridine, afforded the title
10 compound with 18 % overall yield.

¹H NMR (CDCl₃, 500 MHz): 7.90 (1H, m), 7.49 (1H, m), 7.40 (1H, m), 6.81 (2H, m), 6.67 (2H, m), 3.12 (4H, m), 2.77 (2H, q, *J* = 7.56 Hz), 2.78 (3H, s), 2.58 (4H, m), 2.36 (3H, s), 2.35 (3H, s), 1.15 (3H, t, *J* = 7.56 Hz); MS (ESI⁺ TOF): 375 (M⁺)

15

EXAMPLE 8

(3-Ethyl-6-methoxy-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure outlined in Step 4 of Example 1, but substituting
20 4-chloro-3-ethyl-6-methoxy-2-methylquinoline for 9-chloroacridine, afforded the title compound with 5 % overall yield.

MS (ESI⁺ TOF): 391 (M⁺)

EXAMPLE 9

25 **(3-Ethyl-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine**

Following the procedure outlined in Step 4 of Example 1, but substituting 4-chloro-3-ethyl-2-methylquinoline for 9-chloroacridine, afforded the title compound with 17 % overall yield.

25

¹H NMR (CDCl₃, 500 MHz): 7.91 (1H, m), 7.72 (1H, m), 7.55 (1H, m), 7.25 (1H, m), 6.80 (2H, m), 6.67 (2H, m), 5.72 (1H, s), 3.11 (4H, m), 2.79 (2H, q, *J* = 7.59 Hz), 2.77 (3H, s), 2.57 (4H, m), 2.34 (3H, s), 1.17 (3H, t, *J* = 7.59 Hz); MS (ESI⁺ TOF): 361 (M⁺)

5

EXAMPLE 10

(4-Methoxyacridin-9-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure outlined in Step 3 of Example 1, but substituting 9-hydroxy-4-methoxyacridine for 9-(10H)acridone, afforded 9-chloro-4-methoxyacridine, which was reacted with 4-(4-methylpiperazin-1-yl)aniline (according to the Step 4 in the procedure of Example 1) to obtain the title compound with 24 % overall yield.

MS (ESI⁺ TOF): 399 (M⁺)

15

EXAMPLE 11

[4-(4-Methylpiperazin-1-yl)phenyl]-(1,2,3,4-tetrahydroacridin-9-yl)amine

Following the procedure outlined in Step 3 of Example 1, but substituting 1,2,3,4-tetrahydro-9-(10H)acridone for 9-(10H)acridone, afforded 9-chloro-1,2,3,4-tetrahydroacridine, which was reacted with 4-(4-methylpiperazin-1-yl)aniline (according to Step 4 in the procedure of Example 1) to obtain the title compound with 46 % overall yield.

¹H NMR (CDCl₃, 500 MHz): 7.96 (1H, m), 7.72 (1H, m), 7.56 (1H, m), 7.27 (1H, m), 6.83 (2H, m), 6.73 (2H, m), 5.82 (1H, s), 3.14 (6H, m), 2.67 (2H, m), 2.58 (4H, m), 2.35 (3H, s), 1.94 (2H, m), 1.84 (2H, m); MS (ESI⁺ TOF): 373 (M⁺)

25

EXAMPLE 12

Acridin-9-yl-[4-(piperidin-1-yl)phenyl]amine

0.202 g (1.0 mmol) of 1-bromo-4-nitrobenzene and 0.20 ml of piperidine (2.0 mmol) were dissolved in 3 ml of dimethylsulfoxide. 0.207 g (1.5 mmol) of potassium carbonate was added and the reaction mixture was heated to 100°C. After 2 h 100 ml of water was added, and the mixture was extracted a few times with dichloromethane. The combined organic layers were dried over sodium sulfate and evaporated to obtain crude 4-(piperidin-1-yl)-1-nitrobenzene. Following the procedure outlined in Step 2 of Example 1, but substituting 4-piperidin-1-yl-1-nitrobenzene for 1-methyl-4-(4-nitrophenyl)piperazine, afforded 4-piperidin-1-ylaniline, which was reacted with 9-chloroacridine (according to Step 4 in the procedure of Example 1) to obtain the title compound with 6 % overall yield.

¹H NMR (CDCl₃, 500 MHz): 8.05 (2H, m), 8.00 (2H, m), 7.48 (2H, m), 7.09 (4H, m), 6.86 (2H, m), 3.13 (4H, m), 1.71 (4H, m), 1.58 (2H, m); MS (ESI⁺ TOF): 354 (M⁺)

EXAMPLE 13

Acridin-9-yl-[4-(4-methylpiperidin-1-yl)phenyl]amine

Following the procedure of Example 12, but substituting 4-methylpiperidine for piperidine, afforded the title compound with 6 % overall yield.

MS (ESI⁺ TOF): 368 (M⁺)

EXAMPLE 14

Acridin-9-yl-[4-(3-hydroxymethylpiperidin-1-yl)phenyl]amine

Following the procedure of Example 12, but substituting 3-hydroxymethylpiperidine for piperidine, afforded the title compound with 3 % overall yield.

27

¹H NMR (CDCl₃, 500 MHz): 7.93 (2H, m), 7.81 (2H, m), 7.47 (2H, m), 7.04 (4H, m), 6.92 (2H, m), 3.68 (2H, m), 3.59 (2H, m), 2.78 (1H, m), 2.61 (1H, m), 1.95 (1H, br s), 1.83 (2H, m), 1.72 (1H, m), 1.23 (2H, m); MS (ESI⁺ TOF): 384 (M⁺)

5

EXAMPLE 15**Acridin-9-yl-[4-(pyrrolidin-1-yl)phenyl]amine**

Following the procedure of Example 12, but substituting pyrrolidine for piperidine, afforded the title compound with 50 % overall yield.

10 ¹H NMR (CDCl₃, 500 MHz): 8.00 (4H, m), 7.56 (2H, m), 7.18 (2H, m), 7.00 (2H, m), 6.50 (2H, m), 3.27 (4H, m), 2.02 (4H, m); MS (ESI⁺ TOF): 340 (M⁺)

EXAMPLE 16

15 **Acridin-9-yl-[4-(piperazine-1-yl)phenyl]amine**

Following the procedure outlined in Step 2 of Example 1, but substituting N-(4-nitrophenyl)piperazine for 1-methyl-4-(4-nitrophenyl)piperazine, afforded 4-(piperazin-1-yl)aniline, which was reacted with 9-chloroacridine (according to Step 4 in the procedure of Example 1) to obtain the title compound with 16 %
20 overall yield.

¹H NMR (CD₃OD, 500 MHz): 8.21 (2H, m), 7.94 (4H, m), 7.42 (2H, m), 7.38 (2H, m), 7.20 (2H, m), 3.56 (4H, m), 3.43 (4H, m); MS (ESI⁺ TOF): 355 (M⁺)

25

EXAMPLE 17**Acridin-9-yl-[4-(4-acetylpiperazine-1-yl)phenyl]amine**

35 mg (0.10 mmol) of 9-[4-(piperazin-1-yl)phenyl]aminoacridine (Example 16) was dissolved in 2 ml of chloroform, 7.1 μl (0.10 mmol) of acetyl chloride and catalytical amounts of pyridine and triethylamine were added. After

2 h of stirring at room temperature, the solvents were removed under vacuum, and the residue was purified by silica gel chromatography (eluent chloroform : methanol 6:1). The fractions with the title compound were combined, evaporated, taken up to water and extracted with chloroform. After drying over sodium sulfate, the organic layer was evaporated to obtain the title compound (6 %, overall yield 1 %).

MS (ESI⁺ TOF): 397 (M⁺)

EXAMPLE 18

10 **Acridin-9-yl-[4-(4-benzylpiperazine-1-yl)phenyl]amine**

Following the procedure outlined in Step 1 of Example 1, but substituting 9-[4-(piperazin-1-yl)phenyl]aminoacridine (Example 16) for N-(4-nitrophenyl)piperazine, and benzyl bromide for methyl iodide, afforded the title compound (15 %, overall yield 2 %).

15 ¹H NMR (CDCl₃, 500 MHz): 8.00 (4H, m), 7.48 (2H, m), 7.34 (4H, m), 7.28 (1H, m), 7.12 (4H, m), 6.88 (2H, m), 3.59 (2H, s), 3.21 (4H, m), 2.63 (4H, m); MS (ESI⁺ TOF): 445 (M⁺)

EXAMPLE 19

20 **Acridin-9-yl-[4-(4-isopropylpiperazine-1-yl)phenyl]amine**

Following the procedure outlined in Step 1 of Example 1, but substituting 9-[4-(piperazin-1-yl)phenyl]aminoacridine (Example 16) for N-(4-nitrophenyl)piperazine, and isopropyl iodide for methyl iodide, afforded the title compound (11 %, overall yield 2 %).

25 MS (ESI⁺ TOF): 397 (M⁺)

EXAMPLE 20

2-{4-[4-(1,2,3,4-Tetrahydroacridin-9-yl)aminophenyl]piperazin-1-yl}ethanol

0.14 ml (2.0 mmol) of 2-bromoethanol and 0.17 ml (2.4 mmol) of acetyl chloride were dissolved in 2 ml of dichloromethane. 0.28 ml (2.0 mmol) of triethylamine was added, and the reaction mixture was stirred at room temperature. After 1 h 50 ml of dichloromethane was added to the reaction mixture, which was then washed with a sodium bicarbonate (sat.) solution, a 10 % citric acid solution and water. The organic phase was dried over sodium sulfate and evaporated to obtain 0.241 g (72 %) of 2-bromoethylacetate. Following the procedure outlined in Step 1 of Example 1, but substituting 2-bromoethylacetate for methyl iodide, afforded 2-[4-(4-nitrophenyl)piperazin-1-yl]ethylacetate (64 %), which was reduced according to the procedure of Step 2 in Example 1 to give the corresponding amine 2-[4-(4-aminophenyl)piperazin-1-yl]ethylacetate in quantitative yield. 2-[4-(4-aminophenyl)piperazin-1-yl]ethylacetate was reacted with 9-chloro-1,2,3,4-tetrahydroacridine (according to the procedure of Step 4 in Example 1) to obtain 2-{4-[4-amino-(1,2,3,4-tetrahydroacridin-9-yl)phenyl]piperazin-1-yl}ethyl acetate with 47 % yield. Treating the ester with 4 equivalents of lithium hydroxide in a dioxane /water solution overnight, afforded the title compound with 60 % yield (overall yield 20 %).

MS (ESI⁺ TOF): 403 (M⁺)

EXAMPLE 21

2-{4-[4-(Acridin-9-yl)aminophenyl]piperazin-1-yl}ethanol

Following the procedure outlined in Step 1 of Example 1, but substituting 9-[4-(piperazin-1-yl)phenyl]aminoacridine (Example 16) for N-(4-nitrophenyl)piperazine, and 2-bromoethylacetate for methyl iodide, afforded 2-{4-[4-amino-(1,2,3,4-tetrahydroacridin-9-yl)phenyl]piperazin-1-yl}ethyl acetate (31 %). Treating the ester with 8 equivalents of lithium hydroxide in dioxane /water solution overnight, afforded the title compound with 37 % yield (overall yield 2 %).

MS (ESI⁺ TOF): 399 (M⁺)

EXAMPLE 22

Acridin-9-yl-[4-(4-cyclopropylpiperazine-1-yl)phenyl]amine

35 mg (0.10 mmol) of 9-[4-(piperazin-1-yl)phenyl]aminoacridine (Example 16) was dissolved in 1 ml of methanol. 57 μ l (1.0 mmol) of acetic acid, 22 μ l (0.11 mmol) of (1-ethoxycyclopropyloxy)-trimethylsilane and a small amount of 3 Å molecular sieves were added. The reaction mixture was stirred at room temperature under nitrogen atmosphere. After 30 min 28 mg (0.45 mmol) of sodium cyanoborohydride was added and the reaction mixture was heated at 50°C over night. The solvents were removed under vacuum, and the residue was purified by chromatography (silica gel column, eluent chloroform/methanol 6:1) to obtain 3.8 mg (10 %, overall yield 2 %) of the title compound.

MS (ESI⁺ TOF): 395 (M⁺)

15 EXAMPLE 23

4-[4-(4-Methylpiperazin-1-yl)phenylamino]quinoline-3-carbonitrile

0.91 ml (10 mmol) of aniline and 1.69 g (10 mmol) of ethyl(ethoxymethylene)cyanoacetate were dissolved in 10 ml of pyridine and heated under reflux. After 3 h pyridine was removed under vacuum, and the residue was purified by chromatography (silica gel column, eluent 1 % methanol in dichloromethane). 1.08 g (50 %) of ethyl(anilinomethylene)cyanoacetate were obtained. The compound was cyclized by heating in a biphenyl/phenyl ether mixture. After cooling a precipitate was filtered and washed with diethyl ether to give 4-hydroxyquinoline-3-carbonitrile (49 %). Following the procedure outlined in Step 3 of Example 1, but substituting 4-hydroxyquinoline-3-carbonitrile for 9-(10H)acridone, 4-chloro-3-cyanoquinoline (90 %) was obtained, which was reacted with 4-(4-methylpiperazin-1-yl)aniline (according to the procedure of Step 4 in Example 1), to afford the title compound with 12 % yield (overall yield 3 %).

¹H NMR (CDCl₃, 500 MHz): 8.66 (1H, s), 8.01 (1H, m), 7.78 (1H, m), 7.73 (1H, m), 7.42 (1H, m), 7.15 (2H, m), 6.94 (2H, m), 3.29 (4H, m), 2.63 (4H, m), 2.39 (3H, s); MS (ESI⁺ TOF): 344 (M⁺)

EXAMPLE 24

(3-Isopropyl-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

4.56 ml (50 mmol) of aniline and 10.7 ml (60 mmol) of ethyl 2-isopropylacetoacetate were mixed with 50 ml of chloroform. 0.48 g (2.5 mmol) of *para*-toluenesulfonic acid was added, and the reaction mixture was refluxed with continuous removal of the water produced in the reaction. After 2 d chloroform was removed in vacuum, and the residue was refluxed in 10 ml of phenyl ether. After cooling, the precipitate was filtered and washed with diethyl ether to give 4-hydroxy-3-isopropyl-2-methylquinoline (21 %). Following the procedure of Step 3 in Example 1, but substituting 4-hydroxy-3-isopropyl-2-methylquinoline for 9-(10H)acridone, 4-chloro-3-isopropyl-2-methylquinoline (100 %) was obtained, which was reacted with 4-(4-methylpiperazin-1-yl)aniline (following the procedure of Step 4 in Example 1), to afford the title compound with 61 % yield (overall yield 13 %).

¹H NMR (CDCl₃, 500 MHz): 7.95 (1H, m), 7.73 (1H, m), 7.54 (1H, m), 7.24 (1H, m), 6.79 (2H, m), 6.60 (2H, m), 5.72 (1H, s), 3.61 (1H, q, *J* = 7.28), 3.11 (4H, m), 2.81 (3H, s), 2.57 (4H, m), 2.34 (3H, s), 1.38 (6H, d, *J* = 7.28); MS (ESI⁺ TOF): 375 (M⁺)

20

EXAMPLE 25

(2,3-Dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure of Example 24, but substituting ethyl 2-methylacetoacetate for ethyl 2-isopropylacetoacetate, afforded the title compound with 11 % yield.

25

¹H NMR (CDCl₃, 500 MHz): 8.02 (1H, m), 7.78 (1H, m), 7.59 (1H, m), 7.33 (1H, m), 6.83 (2H, m), 6.69 (2H, m), 5.90 (1H, s), 3.13 (4H, m), 2.73 (3H, s), 2.58 (4H, m), 2.35 (3H, s), 2.24 (3H, s); MS (ESI⁺ TOF): 347 (M⁺)

EXAMPLE 26

[4-(4-Methylpiperazin-1-yl)phenyl]-(2-methyl-1,2,3,4-tetrahydro-acridin-9-yl)amine

1.37 g (10 mmol) of 2-aminobenzoic acid and 1.23 ml (10 mmol) of 4-methylcyclohexanone were dissolved in 10 ml of phosphorus oxychloride and the reaction mixture was heated under reflux in a nitrogen atmosphere. After 3 h most of the phosphorus oxychloride was removed under vacuum. The remaining brown syrup was poured into a cold sodium bicarbonate (sat.) solution and washed once with chloroform. The yellow precipitate forming in the basic water solution was filtered (2-methyl-1,2,3,4-tetrahydro-9-(10H)-acridone, 46 %). Following the procedure of Step 3 in Example 1, but substituting 2-methyl-1,2,3,4-tetrahydro-9-(10H)acridone for 9-(10H)acridone, gave 9-chloro-2-methyl-1,2,3,4-tetrahydroacridine (40 %), which was reacted with 4-(4-methylpiperazin-1-yl)aniline (according to the procedure of Step 4 in Example 1), to afford the title compound with 39 % yield (overall yield 7 %).

¹H NMR (CDCl₃, 500 MHz): 7.97 (1H, m), 7.70 (1H, m), 7.55 (1H, m), 7.24 (1H, m), 6.83 (2H, m), 6.72 (2H, m), 5.81 (1H, br s), 3.23 (1H, m), 3.14 (5H, m), 2.86 (1H, m), 2.57 (4H, m), 2.35 (3H, s), 2.24 (1H, m), 2.03 (1H, m), 1.93 (1H, m), 1.57 (1H, m), 1.10 (3H, d); MS (ESI⁺ TOF): 387 (M⁺)

20

EXAMPLE 27

[4-(4-methylpiperazin-1-yl)phenyl]-(7,8,9,10-tetrahydro-6H-cyclohepta[b]quinolin-11-yl)amine

Following the procedure of Example 26, but substituting cycloheptanone for 4-methylcyclo-hexanone, afforded the title compound (0,2 %).

(ESI⁺ TOF): 387 (M⁺)

EXAMPLE 28

[4-(4-Methylpiperazin-1-yl)phenyl]-(2,7-dimethyl-1,2,3,4-tetrahydroacridin-9-yl)amine

Following the procedure of Example 26, but substituting 2-amino-5-methylbenzoic acid for 2-aminobenzoic acid, afforded the title compound (4 %).

¹H NMR (CDCl₃, 500 MHz): 8.10 (1H, m), 7.43 (2H, m), 6.85 (2H, m), 6.83 (2H, m), 6.27 (1H, br s), 3.37 (1H, m), 3.18 (4H, m), 2.78 (1H, m), 2.61 (4H, m), 2.37 (3H, s), 2.33 (3H, s), 2.18 (1H, m), 2.02 (1H, m), 1.91 (1H, m), 1.52 (1H, m), 1.09 (1H, m), 1.10 (3H, d); MS (ESI⁺ TOF): 401 (M⁺)

EXAMPLE 29

(8-Fluoro-1,2,3,4-tetrahydroacridin-9-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure of Example 26, but substituting 2-amino-6-fluorobenzoic acid for 2-aminobenzoic acid, and cyclohexanone for 4-methylcyclohexanone, afforded the title compound (3 %).

¹H NMR (CDCl₃, 500 MHz): 7.89 (1H, m), 7.51 (1H, m), 7.05 (1H, m), 6.87 (2H, m), 6.82 (2H, m), 3.22 (4H, m), 3.14 (2H, m), 2.66 (4H, m), 2.41 (3H, s), 2.34 (2H, m), 1.88 (2H, m), 1.67 (2H, m); MS (ESI⁺ TOF): 391 (M⁺)

EXAMPLE 30

[4-(4-Methylpiperazin-1-yl)phenyl]-(1,1,3,3-tetramethyl-1,2,3,4-tetrahydroacridin-9-yl)amine

1.49 ml (10 mmol) of ethyl 2-aminobenzoate and 1.73 ml (10 mmol) of 3,3,5,5-tetramethyl-cyclohexanone were mixed with 20 ml of toluene. 20 mg (0.1 mmol) of *para*-toluenesulfonic acid was added, and the reaction mixture was refluxed with continuous removal of the water produced in the reaction. After 9 h, toluene was removed under vacuum, and the residue was refluxed in 10 ml of phenyl ether. After cooling, the precipitate was filtered and washed with diethyl ether to obtain 1,1,3,3-tetramethyl-1,2,3,4-tetrahydro-9(10H)-

acridone (18 %). Following the procedure of Step 3 in Example 1, but substituting 1,1,3,3-tetramethyl-1,2,3,4-tetrahydro-9(10H)acridone for 9-(10H)acridone, gave 9-chloro-1,1,3,3-tetramethyl-1,2,3,4-tetrahydroacridine (21 %), which was reacted with 4-(4-methylpiperazin-1-yl)aniline (according to the
5 procedure of Step 4 in Example 1), to afford the title compound (3 %, overall yield 0,1 %).

MS (ESI⁺ TOF): 429 (M⁺)

EXAMPLE 31

10 **(1,4-Methano-1,2,3,4-tetrahydroacridin-9-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine**

Following the procedure of Example 30, but substituting norcamphore for 3,3,5,5-tetramethyl-cyclohexanone, afforded the title compound (overall yield 1 %).

15 ¹H NMR (CDCl₃, 500 MHz): 7.99 (1H, m), 7.84 (1H, m), 7.58 (1H, m), 7.38 (1H, m), 7.02 (2H, m), 6.91 (2H, m), 6.13 (1H, s), 3.44 (1H, m), 3.21 (4H, m), 3.00 (1H, m), 2.61 (4H, m), 2.37 (3H, s), 1.97 (1H, m), 1.77 (2H, m), 1.48 (2H, m), 1.26 (1H, m); MS (ESI⁺ TOF): 385 (M⁺)

20 EXAMPLE 32

[4-(4-Methylpiperazin-1-yl)phenyl]-(1,2,3,4,5,6,7,8-octahydroacridin-9-yl)amine

Following the procedure of Example 30, but substituting ethyl 2-amino-1-cyclohexene-1-carboxylate for ethyl 2-aminobenzoate, and cyclohexanone for
25 3,3,5,5-tetramethylcyclo-hexanone, afforded the title compound (overall yield 0,2 %).

MS (ESI⁺ TOF): 377 (M⁺)

EXAMPLE 33

(3-Ethyl-2-methylquinolin-4-yl)-methyl-[4-(4-methylpiperazin-1-yl)phenyl]amine

Following the procedure of Step 1 in Example 1, but substituting (3-ethyl-
5 2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine (Example 9) for
N-(4-nitrophenyl)piperazine, afforded the title compound (overall yield 12 %).

¹H NMR (CDCl₃, 500 MHz): 8.03 (1H, m), 7.58 (2H, m), 7.34 (1H, m),
6.81 (2H, m), 6.42 (2H, m), 3.33 (3H, s), 3.07 (4H, m), 2.80 (3H, s), 2.71 (2H, q,
J = 7.50 Hz), 2.57 (4H, m), 2.34 (3H, s), 1.12 (3H, t, J = 7.50 Hz); MS (ESI⁺)
10 TOF): 375 (M⁺)

EXAMPLE 34

Acridin-9-yl-methyl-[4-(4-(methylpiperazin-1-yl)phenyl]amine

Following the procedure of Step 1 in Example 1, but substituting acridin-
15 9-yl-[4-(4-methyl-piperazin-1-yl)-phenyl]-amine (Example 1) for N-(4-
nitrophenyl)piperazine, afforded the title compound (overall yield 9 %).

¹H NMR (CD₃OD, 500 MHz): 7.8 (2H, br s), 7.54 (2H, m), 7.48 (2H, m),
6.98 (4H, m), 6.79 (2H, m), 3.77 (3H, s), 3.18 (4H, m), 2.65 (4H, m), 2.36 (3H,
s); MS (EI⁺): 383 (M⁺)

20

The compounds of the invention show interesting pharmacological
properties, namely, they exhibit antagonistic affinity for alpha-2 adrenoceptors.
This activity is demonstrated in the pharmacological tests presented below.

25 **EXPERIMENT I: Binding affinity**

The affinity of test compounds for the three human alpha-2-adrenoceptor
subtypes (alpha-2A, alpha-2B and alpha-2C) was determined in binding
competition assays with ³H-rauwolscine. The biological material consisted of
membranes from Shionogi S115 cells stably transfected with one of the three

- human alpha-2 subtypes (A. Marjamäki et al., Biochem. Biophys. Acta, vol.1134, 1992, p.169). The membrane suspension (about 10 µg total protein per sample) and 1 nM of ³H-rauwolscine (specific activity 75-85 Ci/mmol) were incubated with a minimum of six concentrations of the test compound in a total volume of 90 µl (50 mM KH₂PO₄, pH 7.5, at room temperature). Non-specific binding was defined by 100 µM oxymetazoline and corresponded to 4-10 % of the total binding. After 30 min at room temperature, the incubation was terminated by rapid filtration (TomTec 96 harvester) through presoaked GF/B glass-fiber mats (Wallac Oy) and three washes with ice-cold 50 mM KH₂PO₄ (pH 7.5, at room temperature). After drying, a solid scintillate (Meltilex; Wallac Oy) was melted on filter mats, and the radioactivity was measured (BetaPlate; Wallac Oy). The analysis of the experiments was carried out by non-linear least square curve fitting. IC₅₀'s were converted to K_i's by using the equation of Cheng-Prusoff
- ($K_i = IC_{50} / (1 + [^3H\text{-ligand}] / K_d, ^3H\text{-ligand})$). The K_i values for Compound 1 obtained in a minimum of three independent experiments were:

alpha-2A Adrenoceptor: 3150 ± 50 nM

alpha-2B Adrenoceptor: 1470 ± 130 nM

alpha-2C Adrenoceptor: 28 ± 2 nM

20

EXPERIMENT II: Antagonist activity

- Antagonist activity was determined as the ability of compounds to competitively inhibit epinephrine-stimulated ³⁵S-GTPγS binding to G proteins (J.R. Jasper et al., Biochem. Pharmacol., vol.55, 1998, p.1035) in membranes of CHO cells stably transfected with one of the three human alpha-2 subtypes (K. Pohjanoksa et al., Eur. J. Pharmacol., vol.335, 1997, p.53). Membranes (5-10 µg of protein per sample) and 12 concentrations of test compound were preincubated for 30 min at room temperature in 50 mM Tris, 5 mM MgCl₂, 150 mM NaCl, 1 mM DTT, 1 mM EDTA, 10 µM GDP, 30 µM ascorbic acid, pH 7.4, with a fixed concentration of epinephrine (5 µM for alpha-2A, 15 µM for alpha-2B, 5 µM for alpha-2C). Then trace amounts of ³⁵S-GTPγS (0.08 nM- 0.15 nM, specific activity 1250 Ci/mmol) were added to the incubation mixture. After an additional 30 min at room temperature, the incubation was terminated by rapid vacuum filtration through glass fiber filter. Filters were washed three times with

30

37

5 ml ice cold wash buffer (20 mM Tris, 5 mM MgCl₂, 1 mM EDTA, pH 7.4, at room temperature), dried and counted for radioactivity in a scintillation counter. Analysis of experiments was carried out by nonlinear least square fitting. Experiments were repeated at least three times. The K_B values of Compound 1 were found to correspond to:

alpha-2A Adrenoceptor: 1495 ± 270 nM

alpha-2B Adrenoceptor: 2175 ± 345 nM

alpha-2C Adrenoceptor: 16 ± 6 nM

10 **EXPERIMENT III: Antagonism of dexmedetomidine –induced locomotor inhibition by atipamezole but not by Compound 1; demonstration of *in vivo* alpha2C selectivity of Compound 1**

Dexmedetomidine and atipamezole are very potent and specific alpha-2 adrenoceptor agonists and antagonists, respectively, which lack alpha-2 subtype selectivity (H. Scheinin et al., European Journal of Pharmacology, Molecular Section, vol.151(1), 1988, p.35-42). The alpha-2 agonist –induced sedation is known to be an alpha-2A –mediated phenomenon that can be antagonised by alpha-2 antagonists (J. Sallinen et al., Mol. Pharmacol. Vol.51, 1997, p.36-46, and A. Haapalinna et al., Naunyn-Schmiedeberg's Arch. Pharmacol. Vol.356, 1997, p.570-582). The sedative effect of alpha-2 agonists in mice is measured by the inhibition on locomotor activity. Therefore, we compared the ability of Compound 1 and atipamezole to antagonise the dexmedetomidine –induced locomotor inhibition to evaluate the *in vivo* alpha-2A adrenoceptor antagonism (and alpha-2C selectivity) of these compounds.

25 Spontaneous locomotor activity of a total of 76 male NMRI mice (B&K, Sweden) was measured by placing individual animals into a polypropylene animal cage (38 x 22 x 15 cm). The cages were surrounded with an infrared photobeam frame system designed for activity measurements (Photobeam Activity System PAS, Cage Rack, San Diego Instruments, San Diego, CA, USA). The animals were injected with various doses of either Compound 1 or atipamezole 20 min before injection of dexmedetomidine (50 nmol/kg s.c.). Spontaneous locomotor activity was measured 20 min after dexmedetomidine injection.

The results presented in figure 1 show that atipamezole inhibited dexmedetomidine –induced sedation with doses 0.3 and 1.0 $\mu\text{mol/kg}$ s.c. ($p < 0.01$) as was expected. In contrast, Compound 1 did not antagonise the alpha-2-agonist –induced sedation at all, demonstrating the lack of alpha-2A antagonism and the alpha-2C selectivity of Compound 1 *in vivo*.

EXPERIMENT IV: Stress-protective effect of Compound 1 in the mouse forced swimming test

Exposure of a test animal to an intense stressful stimulus has been observed to propagate a state of behavioural despair. This can be observed for example employing the forced swimming test, in which a rat or mouse is put into a water-filled cylinder. After a vigorous period of attempts to escape animals adopt an immobile floating posture; the extent of the immobile period is monitored and can be reduced by antidepressants and stress-protective agents. Transgenic mice that lack functional alpha-2C adrenoceptor tolerated swim-stress better than their similarly treated wild-type controls (J. Sallinen et al., Mol. Psychiatry, vol.4, 1999, p.443-452). Therefore, an increase in the forced swimming activity can be used as a measure of *in vivo* alpha-2C selective antagonism of a compound. The non-selective antagonist atipamezole did not have a clear stress-protective effect and even increased vocalizations of test animals (T. Kauppila et al., Eur. J. Pharmacol., vol.205, 1991, p.177-182). This may have resulted from the simultaneous alpha-2A antagonistic activity of atipamezole, since conventionally non-selective alpha-2 adrenoceptor antagonists, such as yohimbine, have been found to be anxiogenic (S. Southwick et al., Arch. Gen. Psychiatry, vol.54, 1997, p.749-758).

The forced swim test was conducted as originally described employing the stress sensitive (J. Crawley and L. Davis, Brain Research Bulletin, vol.8, 1982, p.609-612, and US-A-5 902 807) Balb/c mouse strain (B&K, Sweden). The mice were administered either with vehicle (0.1% DMSO, 5 ml/kg subcutaneously), Compound 1 0.3 $\mu\text{mol/kg}$, or atipamezole 0.3 $\mu\text{mol/kg}$ 40 min before putting the mice into a vessel (10 cm diameter, 18.5 cm height, filled with 25°C water up to 8 cm height). The cumulative activity of each mouse was measured between 2 and 6 min after introduction to the vessel. Only vigorous

attempts to escape (climbing) were registered. The mice (a total of 96) were tested only once. The results are shown in Figure 2.

Mice administered with Compound 1 tolerated clearly better stress-induced propagation of behavioural despair when compared to vehicle injected group of mice (1-way ANOVA, followed by LSD post hoc test; $p = 0.013$) as was expected for an alpha-2C selective compound. Atipamezole did not have any clear effect on the activity compared to control mice ($p = 0.52$). A possible marginal effect of atipamezole was expected, since this subtype-non-selective alpha-2 antagonist blocks also the alpha-2C adrenoceptors. On the other hand the employed 0.3 $\mu\text{mol/kg}$ dose of atipamezole was shown to have also alpha2A antagonism *in vivo* (Figure 1). The employed doses of atipamezole have also been shown to have clear neurochemical effects, i.e. to stimulate noradrenaline release in brain (A. Haapalinna et al., Naunyn-Schmiedeberg's Arch. Pharmacol. Vol.356, 1997, p.570-582). Therefore, the results support that alpha-2A adrenoceptor antagonism of atipamezole can counteract the stress-protective and antidepressant effects of alpha-2C antagonism *in vivo* (J. Sallinen et al., Mol. Psychiatry, vol.4, 1999, p.443-452, and US-A-5 902 807).

In general, the compounds of the invention exhibiting alpha-2 adrenoceptor antagonistic activity may be useful for the treatment of diseases or conditions wherein alpha-2 antagonists are effective. For example, the compounds can be used to treat disorders of the central nervous system, male sexual impotence, orthostatic hypotension, non-insulin dependent diabetes, and obesity, for example, disorders of the central nervous system. The compounds can also be used to reverse effects induced by alpha-2 agonists. Disorders of the central nervous system treatable with the compounds of the invention include depression, anxiety, post traumatic stress disorder, schizophrenia, Parkinson's disease, and other movement disorders.

The selective alpha-2C antagonists of the present invention may be used for the treatment of various diseases or conditions of CNS-system where alpha2C-antagonists are indicated to be beneficial (see e.g. US-A-5 902 807, J. Sallinen et al., Neuroscience, vol. 86, 1998, p.959-965, J. Sallinen et al., J. Neurosci., vol.18, 1998, p.3035-3042, and M. Bjorklund et al., Molecular Pharmacology, vol.54, 1998, p.569-76, the contents of which are hereby incorporated by reference), for example, in the treatment of schizophrenia and depression. Furthermore, the present alpha-2C antagonists can be used as stress-protective agents, or as agents for the treatment of CNS-disorders

induced by stress, e.g. of post-traumatic stress disorder, as indicated, for example, in US-A-5 902 807 cited above. Because alpha-2C antagonists appear to stimulate central dopaminergic activity, they can be used as antiparkinsonian agents in Parkinson's disease and other movement disorders.

5 Moreover, the present alpha-2C antagonists may also exhibit cognition enhancing properties and thus may be used in the treatment of Alzheimer's disease and other dementias.

Due to the selectivity of tissue distribution, the alpha-2C antagonists of the invention have less or no undesirable side-effects, such as cardiovascular effects.

10

The compounds of the invention may be administered enterally, topically or parenterally.

The compounds of the invention may be formulated alone or together with another active ingredient and/or together with a pharmaceutically acceptable diluent, carrier and/or excipient in different pharmaceutical unit dosage forms, e.g. tablets, capsules, solutions, emulsions and powders etc., depending on the route of administration, using conventional techniques. The pharmaceutically acceptable diluent, carrier and/or excipient can be selected from those conventionally used in the field of pharmaceuticals noticing the chosen route of administration.

15

20

The amount of the active ingredient in a dosage form may vary from, for example, 0.01 to 75 weight-% depending on, for example, the type of the dosage form.

The specific dose level of the compounds of the invention depends on several factors such as the compound to be administered, the species, age and the sex of the subject to be treated, the condition to be treated and on the route and method of administration. Accordingly, the dosage for parenteral administration is typically from 0.5 µg/kg to 10 mg/kg per day and that for oral administration is from 5 µg/kg to 100 mg/kg for an adult male.

25

The present invention further provides a compound of the invention for use as alpha-2 antagonist. Furthermore, a method for the treatment of diseases or conditions where alpha-2 antagonists, e.g. alpha-2C antagonists, are indicated to be useful, e.g. a method for the treatment of diseases or conditions of the central nervous system, is provided. In such a method a

30


therapeutically effective amount of a compound of the invention is administered to a subject in need of such treatment. The use of the compounds of the invention for the manufacture of a medicament to be used for the above indications is also provided.

- 5 Those skilled in the art will appreciate that the embodiments described in this application could be modified without departing from the broad inventive concept. Those skilled in the art also understand that the invention is not limited to the particular disclosed embodiments, but is intended to also cover modifications to the embodiments that are within the spirit and scope of the
- 10 invention.

CLAIMS

1. Use of a compound of formula I,

5



wherein,

R₁ is H or (C₁-C₆)alkyl;

10

each R₂ is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or hydroxy(C₁-C₆)alkyl;

15

A is a benzene ring or (C₅-C₇)cycloalkyl;


when A is a benzene ring each R₃ is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbonyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

20

when A is (C₅-C₇)cycloalkyl each R₃ is independently OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;

43


R₄ and R₅ form, together with the N-atom to which they are attached,



wherein X is O or =NR₆; R₆ is H, OH, NH₂, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, CN-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-, NH₂-CO-,
 5 mono- or di(C₁-C₆)alkylcarbamoyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, phenyl, naphthyl or benzyl, wherein the said phenyl, naphthyl or benzyl is optionally substituted with one to three substituent(s) each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or halo-(C₁-C₆)alkyl;

10

or R₄ and R₅ form, together with the N-atom to which they are attached,



wherein n = 1 or 2; R₆ is as defined above; and r = 0 to 3;

15

or R₄ and R₅ form, together with the N-atom to which they are attached, 1-imidazolyl, 1-imidazolyl or 1-triazolyl, each of which can optionally be substituted with one to three substituent(s) R₇ each independently being (C₁-C₆)alkyl and NH₂;

20

or one of R₄ and R₅ is -SO₂R₈ and the other of R₄ and R₅ is H or (C₁-C₆)alkyl; R₈ is (C₁-C₆)alkyl, phenyl, naphthyl or benzyl, wherein the said phenyl, naphthyl or benzyl is optionally substituted with one to three substituent(s) R₉ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino;

25

R_a and R_b are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed benzene ring optionally substituted with one to three substituent(s) R₃ each independently being OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)-alkylcarbamoyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally substituted with one to four substituent(s) R₁₀ each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed bicyclo[2.2.1]-heptane ring optionally substituted with one to four substituent(s) each independently being OH, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five or six membered heterocyclic ring with one ring heteroatom =NR₁₁, which heterocyclic ring is optionally substituted with one to three substituent(s) R₁₀ as defined above; R₁₁ is H or (C₁-C₆)alkyl, or R₁₁ is phenyl optionally substituted with one to three substituents R₁₂ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino;


m is 0 to 3; and

t is 0 to 3,

or of a pharmaceutically acceptable salt or ester thereof, for the manufacture of a medicament for the treatment of diseases or conditions where antagonists of alpha-2 adrenoceptors are indicated to be useful.

45

2. The use of claim 1, wherein the compound of formula I is a compound of formula IA




or a pharmaceutically acceptable salt or ester thereof,

5

wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} , m and t are as defined in claim 1; i is 1 to 3; and j is 0 to 4.

3. The use of claim 1, wherein the compound of formula I is a compound of formula IB

10



or a pharmaceutically acceptable salt or ester thereof,


15

wherein A , R_1 , R_2 , R_3 , R_4 , R_5 , m and t are as defined in claim 1; and R_a and R_b are independently H, OH, halogen, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, (C_2-C_6) alkynyl, (C_1-C_6) alkoxy, halo (C_1-C_6) alkyl, NO_2 , NH_2 , mono- or di (C_1-C_6) alkylamino, (C_1-C_6) alkyl-S- or CN; or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed five to seven

46

membered carbocyclic ring optionally substituted with one to three substituent(s) R_{10} each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl.

- 5 4. The use of claim 1, wherein the compound of formula I is a compound of formula IC




or a pharmaceutically acceptable salt or ester thereof,

10

wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} , R_{11} , m and t are as defined in claim 1; i is 1 or 2; and j is 0 to 3.

5. The use of claim 1, wherein the compound of formula I is a compound of
15 formula ID



47

or a pharmaceutically acceptable salt or ester thereof,


wherein R_1 , R_2 , R_3 , R'_3 , R_4 , R_5 , m and t are as defined in claim 1; and p is 0 to 3.

5

6. The use of a compound of formula I according to claim 5, wherein m is 1 and R_3 is (C_1-C_6) alkoxy.

7. The use of a compound of formula I according to any one of claims 1 to 6, wherein R_4 and R_5 form, together with the N-atom to which they are attached,

10



wherein X is as defined in claim 1.

8. The use of a compound of formula I according to any one of claims 1 to 7, wherein X is $=NR_6$.

15

9. The use of a compound of formula I according to claim 8, wherein R_6 is (C_1-C_6) alkyl, (C_1-C_6) alkenyl, (C_3-C_6) cycloalkyl or hydroxy (C_1-C_6) alkyl.

20

10. The use of a compound of formula I according to claim 9, wherein R_6 is (C_1-C_6) alkyl.

11. The use of a compound of formula I according to any one of claims 1 to 10 for the manufacture of a medicament for use as a selective α -2C antagonist.


25

12. The use according to any one of claims 1 to 11 for the manufacture of a medicament for the treatment of various disorders of central nervous system.

30

13. A compound of formula II,

48



or of a pharmaceutically acceptable salt or ester thereof,

5 wherein,

R_1 is H or (C₁-C₆)alkyl;


10 each R_2 is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or hydroxy(C₁-C₆)alkyl;

A is a benzene ring or (C₅-C₇)cycloalkyl;

15 when A is a benzene ring each R_3 is independently OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbonyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

20 when A is (C₅-C₇)cycloalkyl each R_3 is independently OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;


R_4 and R_5 form, together with the N-atom to which they are attached,



49

wherein X is O or =NR₆; R₆ is H, OH, NH₂, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, CN-(C₁-C₆)alkyl, (C₁-C₆)alkoxy-CO-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-, NH₂-CO-,
 5 mono- or di(C₁-C₆)alkylcarbamoyl, hydroxy(C₁-C₆)alkyl, (C₃-C₆)cycloalkyl or benzyl, wherein the said benzyl is optionally substituted with one to three
 substituent(s) each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl,
 (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or halo-(C₁-C₆)alkyl;

or R₄ and R₅ form, together with the N-atom to which they are attached,



10 wherein n = 1 or 2; R₆ is as defined above; and r = 0 to 3;

or R₄ and R₅ form, together with the N-atom to which they are attached, 1-imidazolyl, 1-imidazoliny or 1-triazolyl, each of which can optionally be substituted with one to three substituent(s) R₇ each independently being (C₁-
 15 C₆)alkyl or NH₂;

R_a and R_b are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN;

20

or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed benzene ring optionally substituted with one to three substituent(s) R'₃ each independently being OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₁-C₆)alkoxy, halo-(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-
 25 C₆)alkylamino, (C₁-C₆)alkyl-CO-, mono- or di(C₁-C₆)alkylcarbamoyl, (C₁-C₆)alkyl-S-, hydroxy(C₁-C₆)alkyl or NH₂-CO-;

or R_a and R_b form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally
 30 substituted with one to four substituent(s) R₁₀ each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl;

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed bicyclo[2.2.1]-heptane ring optionally substituted with one to four substituent(s) each independently being OH, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

5

or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five or six membered heterocyclic ring with one ring heteroatom =NR₁₁, which heterocyclic ring is optionally substituted with one to three substituent(s) R₁₀ as defined above; R₁₁ is H or (C₁-C₆)alkyl, or R₁₁ is phenyl optionally substituted with one to three substituents R₁₂ each independently being OH, halogen, NO₂, NH₂, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or mono- or di(C₁-C₆)alkylamino;

10

m is 0 to 3; and
t is 0 to 3,

15

with the provisos, that

a) when A is a benzene ring, m is 0 or 1, t is 0, R₁ is H, R₃ is Cl or NO₂, and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed benzene ring, and X is NR₆, then R₆ is not H, -CH₃, -CH₂CH₃, -COCH₃, or -CO-NH₂;

20

b) when A is a benzene ring, then Ra and Rb are not at the same time H;

25

c) when A is a benzene ring, m is 1, t is 0, R₁ is H, and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed benzene ring, which is optionally substituted with Br, and X is O, then R₃ is not NO₂ or -OCH₃;

30

d) when A is a benzene ring, m is 0, t is 0, R₁ is H, and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed nonsubstituted benzene ring, then X is not O;


35

e) the compound is not 4-[4-[(7-Chloro-2-methyl-4-quinolinyl)amino]phenyl]-1-diethylcarbamoylpiperazine, 4-[4-[(6-Chloro-2-methoxy-9-acridinyl)amino]phenyl]-1-diethylcarbamoylpiperazine, 6-amino-4-[[3-chloro-4-(1H-imidazol-1-yl)phenyl]amino]-7-methoxy-3-quinolinecarbonitrile or 4-[[3-

51

chloro-4-(1H-imidazol-1-yl)phenyl]amino]-7-methoxy-6-nitro-3-quinolinecarbonitrile.

14. A compound according to claim 13, which is a compound of formula
5 IIA,



or a pharmaceutically acceptable salt or ester thereof,

- 10 wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} , m and t are as defined in claim 1; i is 1 to 3; and j is 0 to 4.


15 15. A compound according to claim 14, wherein i is 2, j is 0 or 1 and R_{10} is (C_1-C_3) alkyl.

16. A compound according to any one of claims 14 or 15, wherein m is 0 or 1 and R_3 is (C_1-C_3) alkyl or halogen.

17. A compound according to any one of claims 14 to 16, wherein the
20 compound is [4-(4-Methylpiperazin-1-yl)phenyl]-(1,2,3,4-tetrahydroacridin-9-yl)amine, 2-[4-[4-(1,2,3,4-Tetrahydroacridin-9-yl)aminophenyl]piperazin-1-yl]ethanol, [4-(4-Methylpiperazin-1-yl)phenyl]-(2-methyl-1,2,3,4-tetrahydroacridin-9-yl)amine, (8-Fluoro-1,2,3,4-tetrahydroacridin-9-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, [4-(4-Methylpiperazin-1-yl)phenyl]-(2,7-dimethyl-1,2,3,4-tetrahydroacridin-9-yl)amine, [4-(4-Methylpiperazin-1-yl)phenyl]-(7,8,9,10-tetrahydro-6H-cyclohepta[b]quinolin-11-yl)amine or [4-(4-Methylpiperazin-1-yl)phenyl]-(1,1,3,3-tetramethyl-1,2,3,4-tetrahydroacridin-9-yl)amine.
25

52

18. A compound according to claim 13, which is a compound of formula IIB,



5

or a pharmaceutically acceptable salt or ester thereof,

wherein A, R₁, R₂, R₃, R₄, R₅, m and t are as defined in claim 1; and Ra and Rb are independently H, OH, halogen, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, NO₂, NH₂, mono- or di(C₁-C₆)alkylamino, (C₁-C₆)alkyl-S- or CN; or Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring optionally substituted with one to three substituent(s) R₁₀; each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, mono- or di(C₁-C₆)alkylamino or hydroxy(C₁-C₆)alkyl.

15

19. A compound according to claim 18, wherein A is a benzene ring.

20. A compound according to any one of claims 18 or 19, wherein m is 0 or 1.

20

21. A compound according to any one of claims 18 to 20, wherein R₃ is (C₁-C₆)alkyl or (C₁-C₆)alkoxy.

25


22. A compound according to any one of claims 18 to 21, wherein Ra and Rb are independently H or (C₁-C₃)alkyl, wherein the (C₁-C₃)alkyl includes both straight and branched chain radicals of up to 3 carbon atoms.

23. A compound according to claim 18, wherein A is a six membered carbocyclic ring and Ra and Rb form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered carbocyclic ring, wherein said carbocyclic rings can optionally be substituted with one to three substituent(s) each independently being OH, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy or hydroxy(C₁-C₆)alkyl.

24. A compound according to any one of claims 18 to 23, wherein the compound is (3-Ethyl-2,8-dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (2-Methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (3-Ethyl-2,6-dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (3-Ethyl-6-methoxy-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (3-Ethyl-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, 4-[4-(4-Methylpiperazin-1-yl)phenylamino]quinoline-3-carbonitrile, (3-Isopropyl-2-methylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, (2,3-Dimethylquinolin-4-yl)-[4-(4-methylpiperazin-1-yl)phenyl]amine, [4-(4-Methylpiperazin-1-yl)phenyl]-(1,2,3,4,5,6,7,8-octahydroacridin-9-yl)amine or (3-Ethyl-2-methylquinolin-4-yl)-methyl-[4-(4-methylpiperazin-1-yl)phenyl]amine.

20

25. A compound according to claim 13, which is a compound of formula IIC,




25

or a pharmaceutically acceptable salt or ester thereof,

54

wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_{10} , R_{11} , m and t are as defined in claim 1; i is 1 or 2; and j is 0 to 3.

26. A compound according to claim 13, which is a compound of formula
5 IID,



or a pharmaceutically acceptable salt or ester thereof,


wherein R_1 , R_2 , R_3 , R'_3 , R_4 , R_5 , m and t are as defined in claim 1; and p is
10 0 to 3.

27. A compound according to claim 26, wherein m is 1 and R_3 is (C₁-C₆)alkoxy.

15 28. A compound according to any one of claims 26 or 27, wherein the compound is 2-{4-[4-(Acridin-9-yl)aminophenyl]piperazin-1-yl}-ethanol, (4-Methoxyacridin-9-yl)-[4-(4-methylpiperazin-1-yl)-phenyl]amine, Acridin-9-yl-[4-(piperidin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-benzylpiperazin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-methylpiperidin-1-yl)phenyl]amine, Acridin-9-yl-[4-(3-hydroxymethylpiperidin-1-yl)phenyl]amine, Acridin-9-yl-[4-(pyrrolidin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-cyclopropylpiperazin-1-yl)phenyl]amine, Acridin-9-yl-[4-(4-isopropylpiperazin-1-yl)phenyl]amine, (Acridin-9-yl)-methyl-[4-(4-methylpiperazin-1-yl)phenyl]amine or Acridin-9-yl-[2,5-diethoxy-4-(morpholin-4-yl)phenyl]amine.
20

25

29. A compound according to any one of claims 13 to 28, wherein R_4 and R_5 form, together with the N-atom to which they are attached,





wherein R₆ is (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₃-C₆)cycloalkyl or hydroxy(C₁-C₆)alkyl.

5 30. A compound according to claim 29, wherein R₆ is (C₁-C₆)alkyl.

31. A pharmaceutical composition comprising a compound of formula II according to claim 13 as an active ingredient optionally together with a pharmaceutically acceptable diluent, carrier and/or excipient.

THIS PAGE BLANK (USPTO)

1/1

**FIG. 1****FIG. 2**

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number
WO 01/64645 A3

(51) International Patent Classification⁷: C07D 215/44,
219/10, 221/04, 221/06, 221/16, 221/18, 241/04, A61K
31/4706, 31/4709, 31/473, 31/4745, 31/496, A61P 25/00

FIN-20100 Turku (FI). SALO, Harri [FI/FI]: Puolikontie
6 C, FIN-20300 Turku (FI).

(21) International Application Number: PCT/FI01/00203

(74) Agent: ORION CORPORATION; ORION PHARMA,
Industrial Property Rights, P.O.Box 65, FIN-02101 Espoo
(FI).

(22) International Filing Date: 28 February 2001 (28.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
20000480 1 March 2000 (01.03.2000) FI

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicants (*for all designated States except US*): ORION
CORPORATION [FI/FI]: Orionintie 1, FIN-02200 Es-
poo (FI). OY JUVANTIA PHARMA LTD [FI/FI]: Tyk-
istökatu 6 A, BioCity, FIN-20520 Turku (FI).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW). Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR). OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): WURSTER,
Siegfried [DE/FI]: Matarakaari 28 A 1, FIN-21500 Piikkiö
(FI). ENGSTRÖM, Mia [FI/FI]: Sirkkalankatu 12 A 6,
FIN-20500 Turku (FI). SAVOLA, Juha-Matti [FI/FI]:
Siutlankatu 8, FIN-20380 Turku (FI). HÖGLUND, Iisa
[FI/FI]: Kaniikintie 17, FIN-20300 Turku (FI). SALLI-
NEN, Jukka [FI/FI]: Puutarhakatu 24 C 32, FIN-20100
Turku (FI). HAAPALINNA, Antti [FI/FI]: Markulantie
8 A, FIN-20360 Turku (FI). TAUBER, Andrei [RU/FI]:
Väärämäentie 7 d 154, FIN-00700, Helsinki (FI). HOF-
FRÉN, Anna-Marja [FI/FI]: Bilmarkinkatu 4 as 12,


Published:

— with international search report

(88) Date of publication of the international search report:
27 December 2001

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: DERIVATIVES OF QUINOLINE AS ALPHA-2 ANTAGONISTS



(57) Abstract: The invention provides a compound of
formula (I), wherein A, Ra, Rb, R₁ to R₅, m and t are as
defined as disclosed, or a pharmaceutically acceptable salt or
ester thereof, useful as alpha-2 antagonist. The compounds
I can be used for the treatment of diseases or conditions
wherein alpha-2 antagonists are indicated to be effective.

WO 01/64645 A3

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/FI 01/00203

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/44 C07D219/10 C07D221/04 C07D221/06 C07D221/16
 C07D221/18 C07D241/04 A61K31/4706 A61K31/4709 A61K31/473
 A61K31/4745 A61K31/496 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ADRIENNE ADAMS ET AL: "Interaction of DNA-intercalating antitumor agents with adrenoceptors" MOLECULAR PHARMACOLOGY, vol. 27, 1985, pages 480-491, XP002901808 page 487, line 37 - line 40 ---	1-31
X	WO 96 09294 A (WELLCOME FOUNDATION LTD) 28 March 1996 (1996-03-28) abstract; claim 1 ---	1-31
X	GARIMA SATHI ET AL: "New quinolines as potential CNS Agents" ARCH. PHARM., vol. 316, 1983, pages 767-772, XP002901809 Weinheim the whole document ---	1-31
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

12 July 2001

Date of mailing of the international search report

03.09.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Viveca Noren

INTERNATIONAL SEARCH REPORT

International Application No

PLI/FI 01/00203

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>RADZIKOWSKI, CZESLAW ET AL: "New antitumor compounds. Biological studies. IV. Antitumor properties of 21 new acridine derivatives" STN INTERNATIONAL, FILE CAPLUS, CAPLUS ACCESSION NO. 1967:401919, XP002901810 document no. 67:1919 & ARCH. IMMUNOL. THER. EXP. , vol. 15, no. 2, 1967, pages 241-248, ---</p>	1-31
A	<p>KOZYREVA N P ET AL: "Synthesis and study of the anthelmintic activity of some N. heterocycles containing 4-(4'-diethylcarbamoyl-1-'-piperaziny) phenylamine substituents" STN INTERNATIONAL, FILE CAPLUS, CAPLUS ACCESSION NO. 1972:400153, XP002901811 document no. 77:153 & KHIM. -FARM. ZH. , vol. 6, no. 1, 1972, pages 9-12, ---</p>	1-31
A	<p>KLOPMAN GILLES ET AL: "Computer-automated structure evaluation of antileukemic 9-anilinoacridines" STN INTERNATIONAL, FILE CAPLUS, CAPLUS ACCESSION NO. 1987:628423, XP002901812 document no: 107:228423 & MOL. PHARMACOL. , vol. 31, no. 4, 1987, pages 457-476, -----</p>	1-31

INTERNATIONAL SEARCH REPORT

International application No.
PCT/FI 01/00203

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **1-11**
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Claims Nos.: 1-11

Present claims 1-11 concerns the use of the compounds of formula I, as identified in the application, for the manufacture of a medicament for the treatment of diseases of conditions where antagonists of alpha-2-adrenocaptors are indicated to be useful. According to PCT Article 6 the claims should be clear and concise. The wording "diseases or conditons where antagonists of alpha-2 adrenoceptors are indicated to be useful" is not considered to fulfil this requirement as it does not define which diseases and conditions it relates to. The expression "various diseases of central nervous system" in claim 12 is a too wide expression and lacks clarity as the diseases concerned are not defined. It is therefore impossible to perform a complete search of the whole scope claimed. The search has been focused on nervous system diseases in general and alph-2-adrenoceptor antagonists.

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/FI 01/00203

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9609294 A	28-03-1996	AU 3482495 A	09-04-1996
		EP 0782570 A	09-07-1997
		JP 10505600 T	02-06-1998
		TR 960233 A	21-06-1996
		ZA 9507853 A	18-03-1997

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

mis Page Blank (uspto)